



Definition 1. The **Euler Totient Function** is defined for $n \geq 1$ as the number of positive integers not exceeding n which are relatively prime to n .

That is to say:

$$\varphi(n) := \sum_{\substack{k=1 \\ (k,n)=1}}^n 1. \quad (1)$$

Theorem 1. The Euler Totient Function satisfies the equation:

$$\frac{\zeta(s-1)}{\zeta(s)} = \sum_{n=1}^{\infty} \frac{\varphi(n)}{n^s} \quad (2)$$

for $\text{Re}(s) > 1$.

Here $\zeta(s)$ denotes the **Riemann Zeta Function**, defined as

$$\zeta(s) := \sum_{n=1}^{\infty} \frac{1}{n^s} \quad (3)$$

for $\text{Re}(s) > 1$.

Proof. We start by proving that:

$$\varphi(n) = \sum_{d|n} \mu(d) \frac{n}{d} \quad (4)$$

where $\mu(n)$ denotes the **Möbius Function** defined as:

$$\mu(n) = \begin{cases} 1 & \text{if } n = 1; \\ (-1)^k & \text{if } n \text{ is the product of } k \text{ distinct primes;} \\ 0 & \text{if } n \text{ is divisible by a square } > 1. \end{cases} \quad (5)$$

we add some details to the demonstration that can be found in [1]:

Rewrite the definition of $\varphi(n)$ as

$$\varphi(n) = \sum_{k=1}^n \left[\frac{1}{(n, k)} \right] \quad (6)$$

where (n, k) denotes the **Greatest Common Denominator** between n and k . It is a known fact that:

$$\sum_{d|n} \mu(d) = \left[\frac{1}{n} \right]$$

(see [1] for the proof.) Therefore Equation 6 can be written as:

$$\varphi(n) = \sum_{k=1}^n \sum_{\substack{d|(n,k) \\ d|n}} \mu(d) = \sum_{k=1}^n \sum_{\substack{d|n \\ d|k}} \mu(d).$$

Summing over all d that divide both n and k means, for every divisor d of n , summing over all those k in the range $1 \leq k \leq n$ which are multiples of d . If we write $k = qd$ then $1 \leq k \leq n$ if and only if $1 \leq q \leq \frac{n}{d}$. Hence we conclude:

$$\varphi(n) = \sum_{k=1}^n \sum_{\substack{d|n \\ d|k}} \mu(d) = \sum_{d|n} \sum_{q=1}^{\frac{n}{d}} \mu(d) = \sum_{d|n} \mu(d) \sum_{q=1}^{\frac{n}{d}} 1 = \sum_{d|n} \mu(d) \frac{n}{d}.$$

We will use Equation 4 to obtain our main formula:

$$\sum_{n=1}^{\infty} \frac{\varphi(n)}{n^s} = \sum_{n=1}^{\infty} \frac{1}{n^s} \sum_{d|n} \frac{n}{d} \mu(d) = \sum_{n=1}^{\infty} \frac{1}{n^{s-1}} \sum_{d|n} \frac{\mu(d)}{d}. \quad (7)$$

Call $n = dk$ and notice that $d|n$ if and only if $n = dk$, therefore summing over all n is equivalent to summing over all possible d and k , that is to say:

$$\sum_{n=1}^{\infty} \frac{\varphi(n)}{n^s} = \sum_{n=1}^{\infty} \frac{1}{n^{s-1}} \sum_{d|n} \frac{\mu(d)}{d} = \sum_{d=1}^{\infty} \sum_{k=1}^{\infty} \frac{1}{(dk)^{s-1}} \frac{\mu(d)}{d} = \sum_{d=1}^{\infty} \frac{\mu(d)}{d^s} \sum_{k=1}^{\infty} \frac{1}{k^{s-1}}. \quad (8)$$

By definition, for $Re(s) > 1$ we have:

$$\sum_{k=1}^{\infty} \frac{1}{k^{s-1}} = \zeta(s-1)$$

also, it is true that:

$$\sum_{d=1}^{\infty} \frac{\mu(d)}{d^s} = \frac{1}{\zeta(s)}$$

a demonstration of this equation can be found [on our site](#).

Combining these two with equation 8 we conclude:

$$\frac{\zeta(s-1)}{\zeta(s)} = \sum_{n=1}^{\infty} \frac{\varphi(n)}{n^s}.$$

□

References

- [1] Tom M Apostol. *Introduction to analytic number theory*. Springer Science & Business Media, 2013.